The Dahlietta Dahlia Series

Order information – Bosgraaf Greenhouses, Inc. 2012-2013

Minimum Order: Pac Size:	4 trays of 98 cell or 392 plants 98 cell tray – charged for 96 plants
Box size:	21" X 21" X 21" – holds 8 individually boxed boxes
Box weight:	30-35 lbs (6-8 trays)
Availability:	November 6, 2012 – June 10, 2013
Lead time:	8 weeks – late orders DecMay subject to availability of cuttings
Shipped:	FOB Hudsonville, Michigan
Care tags:	Included in price – 1 tag per cutting
Royalty:	Included in price

Handling rooted cuttings – PUSH UP CUTTINGS FROM BOTTOM WITH BOLT

- 1. Unpack immediately and check the need for water.
- 2. Planting dibbling the cutting in pre-watered pots keeps the soil packed around the plant.
- 3. Plant deeply, covering from 1-3 leaves. This helps support the foliage and flowers of the dahlietta.
- 4. A fungicide drench is always a good preventative, especially for pythium.

Handling unrooted cuttings

- 1. Plant cuttings as soon as possible covering the first leaf, or store in a cool place for a very short time period.
- 2. Mist accordingly to conditions during the day with careful monitoring during the first week. Dry down going toward night.
- 3. Maintain bottom heat at 68 F.

5" 1 optional 7-8 week	Container size	# of cuttings	pinching	finish time
10" Hanging Basket 4 pinch 8-9 weel	4" – 4 ½" 5" 6" 7", 7 ½", 8", 8 ½" 10" Hanging Basket	1 1 1 2-3 4	optional optional pinch pinch pinch	6-8 weeks 7-8 weeks 8-9 weeks 8-9 weeks 8-9 weeks 6-8 weeks

<u>Soil</u>

Use a well drained sterilized planting medium, pH of 6.

Fertilization & watering

- 1. Osmocote 14-14-14 can be incorporated into the soil at a rate of 4-6 lbs per cubic yard.
- 2. Constant liquid feed can be used at 150-200 ppm with 20-10-20 or 15-16-17.
- 3. A combination feed using a multi-headed injector with 20-10-20 or 15-0-15 and 20-35 ppm of magnesium.
- 4. Potassium nitrate can be used to build stock stems at 200 ppm.

Temperature & Humidity

Ideal growing temperatures range from 64-70 F. during the day and 65-68 F. at night.

Light conditions

- 1. Grow dahliettas with a maximum amount of available light with a minimum of 2000 footcandles.
- 2. If low light conditions persist, 100 footcandles of HID lighting may be added.

Flowering – Photo Period Requirements

- 1. Long days are needed for quality dahlietta vegetative growth and timely bud development.
- 2. A 14 hour day is required to keep dahliettas vegetative.

Pinching or center bud removal

- 1. Most 4" and 5" productions are not pinched because of great self-branching and early flowering characteristics.
- 2. Center bud removal or first bud removal can be used on non-pinched plants to promote lateral bud/flower development. Flowering is delayed from 4-7 days

Spacing

Allows more air movement and helps prevent fungal and stretching problems

Pest & Disease Prevention

- 1. Thrip, mite, aphid and white fly are potential pests.
- 2. Control can be achieved using Marathon, Conserve, Avid, Maverick, Thiodan, Talstar, Tempo, Orthene, Tame, Decatholon or Duragaard.
- 3. Use of Dursban and Vydate can cause leaf burn.
- 4. A fungicide spray in combination with an insecticide spray is recommended.

Use of growth regulators for height control

- 1. Generally the dahlietta can remain compact.
- 2. Use 5-15 ppm of Bonzi as a spray.

TIPS FOR GROWING DAHLIAS - 2013

1. All dahlias need 14 hours of day length to grow and stay vegetative.

-Growers must extend the day with mum lighting to 14 hours.

-Failure to do so causes dahlias to form tubers, flower profusely, and shut down growth and become dormant.

-Bury 1 set of leaves which helps firm up the plant as it grows.

2. Low temperatures can cause dahlias to grow very little if at all.

-Day and night temperatures of 65 F. keeps dahlias growing.

-We use 68 F. as the benchmark to keep roots moving & vegetation forming.

-Initial temperatures are key to plant growth and development.

-Low temperatures can also cause roots to remain wet and contribute to root rot and other fungal diseases.

3. Using a fungicide drench as a preventive.

-November – February plantings (and later) of dahlia must consider a fungicide planting or within 5 to 7 days.

4. Combating powdery mildew and botrytis.

-Powdery mildew and botrytis need to be considered even when recommended temperatures and conditions are followed

-Changes in the amount of light or swings in daytime temperatures can contribute to conditions causing fungal and mildew problems (humidity, temperature and light levels).

-Use a fungicide spray with an insecticide spray to help with prevention.

- 5. Feed moderately = 100 ppm of 20-10-20, 75 ppm of 15-0-15, 25 ppm mag (200 ppm)
- 6. Height control: most growers are using Bonzi as a spray or drench. Every series is different see color sell sheets for detail.
- 7. Summary: good growing means evaluating plants & conditions daily.

Dahlietta Varieties Available:

- 1. Anna scarlet
- 2. Alexia brilliant red / white edges**
- 3. Betty violet
- 4. Blanca Improved white
- 5. Caroline rose
- 6. Emily light pink
- 7. Julia bright yellow
- 8. Leanne deep burgundy**
- 9. Lily salmon

- 10. Louise rose/white
- 11. Marissa bi-color red/yellow**
- 12. Patricia gold/red
- 13. Patty burgundy
- 14. Rachel orange
- 15. Tessy red
- 16. Wendy pink

**denotes new variety for 2012-2013